

## TRIGONOMETRIJSKE FUNKCIJE DVOSTRUKOG UGLA

Formule su:

1.  $\sin 2\alpha = 2 \sin \alpha \cos \alpha$
2.  $\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha$
3.  $\operatorname{tg} 2\alpha = \frac{2\operatorname{tg} \alpha}{1 - \operatorname{tg}^2 \alpha}$
4.  $\operatorname{ctg} 2\alpha = \frac{\operatorname{ctg}^2 \alpha - 1}{2\operatorname{ctg} \alpha}$

### Primeri:

1) a)  $\sin 2\alpha = \frac{2\operatorname{tg} \alpha}{1 + \operatorname{tg}^2 \alpha}$  **Dokazati.**

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha = (\text{uvek možemo u imenioci dopisati 1, zar ne?})=$$

$$\sin 2\alpha = \frac{2 \sin \alpha \cos \alpha}{1} = \frac{2 \sin \alpha \cos \alpha}{\sin^2 \alpha + \cos^2 \alpha} = (\text{trik: izvučemo zajednički i gore i dole } \cos^2 \alpha)=$$

$$\frac{\cancel{\cos^2 \alpha} \cdot \frac{2 \sin \alpha}{\cos \alpha}}{\cancel{\cos^2 \alpha} \cdot \left( \frac{\sin^2 \alpha}{\cos^2 \alpha} + 1 \right)} = \frac{2 \operatorname{tg} \alpha}{\operatorname{tg}^2 \alpha + 1} = \frac{2 \operatorname{tg} \alpha}{1 + \operatorname{tg}^2 \alpha}$$

b)  $\cos 2\alpha = \frac{1 - \operatorname{tg}^2 \alpha}{1 + \operatorname{tg}^2 \alpha}$  **Dokazati.**

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = \frac{\cos^2 \alpha - \sin^2 \alpha}{1} = \frac{\cos^2 \alpha - \sin^2 \alpha}{\sin^2 \alpha + \cos^2 \alpha} = (\text{isti trik, izvučemo } \cos^2 \alpha \text{ i gore i dole})$$

$$= \frac{\cos^2 \alpha \left( 1 - \frac{\sin^2 \alpha}{\cos^2 \alpha} \right)}{\cos^2 \alpha \left( \frac{\sin^2 \alpha}{\cos^2 \alpha} + 1 \right)} = \frac{1 - \operatorname{tg}^2 \alpha}{\operatorname{tg}^2 \alpha + 1} = \frac{1 - \operatorname{tg}^2 \alpha}{1 + \operatorname{tg}^2 \alpha}, \text{ što je i trebalo dokazati.}$$

v)  $\sin 3\alpha = 3 \sin \alpha - 4 \sin^3 \alpha$  **Dokazati.**

$$\begin{aligned} \sin 3\alpha &= \sin(2\alpha + \alpha) \rightarrow \text{Iskoristimo formulu } \sin(\oplus + \ominus) = \sin\oplus \cos\ominus + \cos\oplus \sin\ominus \\ &= \sin 2\alpha \cos \alpha + \cos 2\alpha \sin \alpha \rightarrow \text{sad formule za dvostruki ugao} \end{aligned}$$

$$\begin{aligned} &= (2 \sin \alpha \cos \alpha) \cos \alpha + (\cos^2 \alpha - \sin^2 \alpha) \cdot \sin \alpha \\ &= 2 \sin \alpha \cos^2 \alpha + \sin \alpha \cos^2 \alpha - \sin^3 \alpha \\ &= 3 \sin \alpha \cos^2 \alpha - \sin^3 \alpha \end{aligned}$$

$$(\text{sad } \acute{\text{c}}emo \text{ iz } \sin^2 \alpha + \cos^2 \alpha = 1 \text{ izraziti } \cos^2 \alpha = 1 - \sin^2 \alpha)$$

$$\begin{aligned} &= 3 \sin \alpha (1 - \sin^2 \alpha) - \sin^3 \alpha \\ &= 3 \sin \alpha - 3 \sin^3 \alpha - \sin^3 \alpha \\ &= 3 \sin \alpha - 4 \sin^3 \alpha \end{aligned}$$

g)  $\cos \alpha = \frac{4}{5}$  **Nadji vrednosti za dvostruke uglove ako je  $\alpha$  u IV kvadrantu.**

Najpre ćemo izračunati  $\sin \alpha$

$$\sin^2 \alpha + \cos^2 \alpha = 1$$

$$\sin^2 \alpha = 1 - \cos^2 \alpha$$

$$\sin^2 \alpha = 1 - \left(\frac{4}{5}\right)^2$$


$$\sin^2 \alpha = 1 - \frac{16}{25}$$

$$\sin^2 \alpha = \frac{9}{25}$$

$$\sin \alpha = \pm \sqrt{\frac{9}{25}}$$

$$\sin \alpha = \pm \frac{3}{5}, \text{ pošto je ugao iz IV kvadranta uzećemo da je } \sin \alpha = -\frac{3}{5}$$

Sada je:



$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$= 2 \left( -\frac{3}{5} \right) \cdot \frac{4}{5}$$

$$= -\frac{24}{25}$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha$$

$$= \left( \frac{4}{5} \right)^2 - \left( -\frac{3}{5} \right)^2 = \frac{16}{25} - \frac{9}{25} = \frac{7}{25}$$

$$\operatorname{tg} 2\alpha = \frac{\sin 2\alpha}{\cos 2\alpha} = \frac{-\frac{24}{25}}{\frac{7}{25}} = -\frac{24}{7}$$

**2) Ako je  $\sin \alpha = 0,6$  i  $\alpha$  pripada prvom kvadrantu, najdi vrednosti za dvostruke uglove.**

Sada ćemo prvo naći  $\cos \alpha$

$$\sin^2 \alpha + \cos^2 \alpha = 1$$

$$\cos^2 \alpha = 1 - \sin^2 \alpha$$

$$\cos^2 \alpha = 1 - (0,6)^2$$

$$\cos^2 \alpha = 1 - 0,36$$

$$\cos^2 \alpha = 0,64$$

$$\cos \alpha = \pm \sqrt{0,64}$$

$$\cos \alpha = \pm 0,8$$

$$\cos \alpha = +0,8$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$= 2 \cdot 0,6 \cdot 0,8$$


$$= 2 \cdot \frac{3}{5} \cdot \frac{4}{5} = \frac{24}{25}$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha$$

$$= \left( \frac{4}{5} \right)^2 - \left( \frac{3}{5} \right)^2 = \frac{7}{25}$$

$$\operatorname{tg} 2\alpha = \frac{\sin 2\alpha}{\cos 2\alpha} = \frac{\frac{24}{25}}{\frac{7}{25}}$$

$$\operatorname{tg} 2\alpha = \frac{24}{7}$$



### 3) Dokazati

$$\text{a) } \sin 15^\circ \cos 15^\circ = \frac{1}{4}$$

$$\begin{aligned} \sin 15^\circ \cos 15^\circ &= (\text{trik je da dodamo } \frac{2}{2}) \\ &= \frac{2 \sin 15^\circ \cos 15^\circ}{2} = (\text{ovo u brojiocu je formula za } \sin 2\alpha = 2 \sin \alpha \cos \alpha) \\ &= \frac{\sin 30^\circ}{2} = \frac{1}{2} = \frac{1}{4} \end{aligned}$$

$$\text{b) } 1 - 4 \sin^2 \alpha \cos^2 \alpha = \cos^2 2\alpha$$

$$\begin{aligned} 1 - 4 \sin^2 \alpha \cos^2 \alpha &= (\text{pošto je formula } \sin 2\alpha = 2 \sin \alpha \cos \alpha, \text{ to je} \\ 4 \sin^2 \alpha \cos^2 \alpha &= \sin^2 2\alpha) \\ \text{pa je } 1 - 4 \sin^2 \alpha \cos^2 \alpha &= 1 - \sin^2 2\alpha = \cos^2 2\alpha \end{aligned}$$

### 4) Dokazati

$$\text{a) } 2 \sin^2 \alpha + \cos 2\alpha = 1$$

$$\begin{aligned} 2 \sin^2 \alpha + \cos 2\alpha &= 2 \sin^2 \alpha + \cos^2 \alpha - \sin^2 \alpha \\ &= \sin^2 \alpha + \cos^2 \alpha = 1 \end{aligned}$$

$$\text{b) } \cos^4 \alpha + \sin^4 \alpha = 1 - 0,5 \sin^2 2\alpha$$

Da bi ovo dokazali podjimo od indentiteta:

$$\sin^2 \alpha + \cos^2 \alpha = 1 / \text{Kvadriramo}$$

$$\sin^4 \alpha + 2 \sin^2 \alpha \cos^2 \alpha + \cos^4 \alpha = 1$$

$$\sin^4 \alpha + \cos^4 \alpha = 1 - 2 \sin^2 \alpha \cos^2 \alpha \quad (\text{dodamo } \frac{2}{2} \text{ izrazu } 2 \sin^2 \alpha \cos^2 \alpha)$$

$$\sin^4 \alpha + \cos^4 \alpha = 1 - \frac{4 \sin^2 \alpha \cos^2 \alpha}{2} \quad (\text{ovde je } 4 \sin^2 \alpha \cos^2 \alpha = \sin^2 2\alpha)$$

$$\sin^4 \alpha + \cos^4 \alpha = 1 - \frac{1}{2} \sin^2 2\alpha$$

$$\sin^4 \alpha + \cos^4 \alpha = 1 - 0,5 \sin^2 2\alpha$$

**5) Dokazati identitet:**

$$\cos 4\alpha + 4\cos 2\alpha + 3 = 8\cos^4 \alpha$$

**Rešenje: Poći ćemo od leve strane da dokažemo desnu.**

$$\begin{aligned} \cos 4\alpha + 4\cos 2\alpha + 3 &= \\ \cos 2 \cdot (2\alpha) + 4(\cos^2 \alpha - \sin^2 \alpha) + 3 &= \\ \cos^2(2\alpha) - \sin^2(2\alpha) + 4\cos^2 \alpha - 4\sin^2 \alpha + 3 &= \\ (\cos^2 \alpha - \sin^2 \alpha)^2 - (2\sin \alpha \cos \alpha)^2 + 4\cos^2 \alpha - 4\sin^2 \alpha + 3 &= \\ (\cos^2 \alpha - (1 - \cos^2 \alpha))^2 - 4\sin^2 \alpha \cos^2 \alpha + 4\cos^2 \alpha - 4\sin^2 \alpha + 3 &= [\text{zamenimo } \sin^2 \alpha = 1 - \cos^2 \alpha] \\ (2\cos^2 \alpha - 1)^2 - 4\cos^2 \alpha(1 - \cos^2 \alpha) + 4\cos^2 \alpha - 4(1 - \cos^2 \alpha) + 3 &= \\ 4\cos^4 \alpha - \cancel{4\cos^2 \alpha} + \cancel{1} - \cancel{4\cos^2 \alpha} + 4\cos^4 \alpha + \cancel{4\cos^2 \alpha} - 4 + \cancel{4\cos^2 \alpha} + 3 &= \\ = 8\cos^4 \alpha & \end{aligned}$$

A ovo smo trebali dokazati!!

**6) Ako je  $\sin \frac{x}{2} + \cos \frac{x}{2} = 1,4$  izračunati  $\sin x$**

**Rešenje:** Kvadriraćemo datu jednakost.

$$\begin{aligned} \sin \frac{x}{2} + \cos \frac{x}{2} &= 1,4 / ()^2 \\ \sin^2 \frac{x}{2} + 2\sin \frac{x}{2} \cos \frac{x}{2} + \cos^2 \frac{x}{2} &= 1,96 \quad [\text{ovde je } 2\sin \frac{x}{2} \cos \frac{x}{2} = \sin x] \\ 1 + \sin x &= 1,96 \\ \sin x &= 1,96 - 1 \\ \sin x &= 0,96 \end{aligned}$$

**7) Predstavi  $\text{tg} 3\alpha$  kao funkciju od  $\text{tg} \alpha$**

**Rešenje:**

$$\begin{aligned} \text{tg} 3\alpha &= \text{tg}(2\alpha + \alpha) = \frac{\text{tg} 2\alpha + \text{tg} \alpha}{1 - \text{tg} 2\alpha \cdot \text{tg} \alpha} = \\ &= \frac{\frac{2\text{tg} \alpha}{1 - \text{tg}^2 \alpha} + \text{tg} \alpha}{1 - \text{tg} \alpha \cdot \frac{2\text{tg} \alpha}{1 - \text{tg}^2 \alpha}} = \frac{\frac{2\text{tg} \alpha + \text{tg} \alpha(1 - \text{tg}^2 \alpha)}{1 - \text{tg}^2 \alpha}}{\frac{1 - \text{tg}^2 \alpha + 2\text{tg}^2 \alpha}{1 - \text{tg}^2 \alpha}} \\ &= \frac{2\text{tg} \alpha + \text{tg} \alpha - \text{tg}^3 \alpha}{1 + \text{tg}^2 \alpha} = \frac{3\text{tg} \alpha - \text{tg}^3 \alpha}{1 + \text{tg}^2 \alpha} \end{aligned}$$

**8) Dokaži indetitet:**

$$\frac{1 + \sin 2\alpha}{\sin \alpha + \cos \alpha} = \sqrt{2} \cos\left(\frac{\pi}{4} - \alpha\right)$$

**Rešenje:**

$$\frac{1 + \sin 2\alpha}{\sin \alpha + \cos \alpha} = \frac{\sin^2 \alpha + \cos^2 \alpha + 2 \sin \alpha \cos \alpha}{\sin \alpha + \cos \alpha} = \frac{(\sin \alpha + \cos \alpha)^2}{\cancel{\sin \alpha + \cos \alpha}}$$

=  $\sin \alpha + \cos \alpha$  = (trik: kod oba sabiraka ćemo dodati  $\frac{2}{2}$  tj.  $\frac{\sqrt{2}^2}{2}$ )

$$\frac{\sqrt{2}^2}{2} \sin \alpha + \frac{\sqrt{2}^2}{2} \cos \alpha = \text{izvučemo } \sqrt{2} \text{ kao zajednički}$$

$$\sqrt{2} \left( \frac{\sqrt{2}}{2} \sin \alpha + \frac{\sqrt{2}}{2} \cos \alpha \right) = \text{pošto je } \sin \frac{\pi}{4} = \frac{\sqrt{2}}{2} \text{ i } \cos \frac{\pi}{4} = \frac{\sqrt{2}}{2} \text{ zamenimo u izraz}$$

$$\sqrt{2} \left( \sin \frac{\pi}{4} \sin \alpha + \cos \frac{\pi}{4} \cos \alpha \right) = \text{malo pretumbamo}$$

$$\sqrt{2} \left( \cos \frac{\pi}{4} \cos \alpha + \sin \alpha \sin \frac{\pi}{4} \right) = \text{ovo u zagradi je formula za } \cos(\alpha - \beta)$$

$$\sqrt{2} \cos\left(\frac{\pi}{4} - \alpha\right)$$